

# CPC COOPERATIVE PATENT CLASSIFICATION

## C CHEMISTRY; METALLURGY

(NOTES omitted)

### CHEMISTRY

## C12 BIOCHEMISTRY; BEER; SPIRITS; WINE; VINEGAR; MICROBIOLOGY; ENZYMOLOGY; MUTATION OR GENETIC ENGINEERING

(NOTES omitted)

**C12P FERMENTATION OR ENZYME-USING PROCESSES TO SYNTHESISE A DESIRED CHEMICAL COMPOUND OR COMPOSITION OR TO SEPARATE OPTICAL ISOMERS FROM A RACEMIC MIXTURE** {(brewing of beer [C12C](#); producing vinegar [C12J](#); producing specific peptides or proteins [C07K](#); producing enzymes [C12N 9/00](#); DNA or RNA concerning genetic engineering, vectors, e.g. plasmids, or their isolation, preparation or purification [C12N 15/00](#); measuring or testing processes involving enzymes or microorganisms [C12Q](#); measuring or testing processes involving nucleic acid amplification reactions [C12Q 1/6844](#); fermentation processes to form a food composition, [A21](#) or [A23](#); compounds in general, see the relevant compound class, e.g. [C01](#), [C07](#))}

#### NOTES

1. This subclass covers the production of compounds or compositions by biochemical transformation of matter performed by using enzymes or microorganisms, wherein microorganisms are defined as any single-celled organisms, including bacteria, fungi, yeast or microalgae, or plant or mammalian cells in the form of cell cultures.
2. In this subclass, documents are primarily classified according to the compounds produced. In addition, if appropriate, classification according to the method or biocatalyst used to produce the compound is made.
3. Classification in groups [C12P 19/14](#) - [C12P 19/24](#), [C12P 39/00](#), [C12P 41/00](#) - [C12P 41/009](#) should only be made together with the corresponding product groups

#### WARNINGS

1. The following IPC groups are not in the CPC scheme. The subject matter for these IPC groups is classified in the following CPC groups:  
 - [C12P 21/04](#) covered by [C07K 7/50](#)  
 - [C12P 21/08](#) covered by [C07K 16/00](#)
2. In this subclass non-limiting references (in the sense of paragraph 39 of the Guide to the IPC) may still be displayed in the scheme.

**1/00 Preparation of compounds or compositions, not provided for in groups [C12P 3/00](#) - [C12P 39/00](#), by using microorganisms or enzymes**

#### NOTES

1. This group is used for the classification of documents relating to the production of compounds of unknown structure
2. When classifying in this group, classification should be made also in [C12R](#)

- 1/02 . by using fungi
- 1/04 . by using bacteria
- 1/06 . by using actinomycetales

**3/00 Preparation of elements or inorganic compounds except carbon dioxide** {(recovery of carbon dioxides as by-products [C12F 3/02](#))}

**5/00 Preparation of hydrocarbons {or halogenated hydrocarbons}**

- 5/002 . {cyclic (compounds containing at least three condensed carbocyclic rings [C12P 15/00](#))}
- 5/005 . . {aromatic (naphthalene [C12P 29/00](#))}
- 5/007 . {containing one or more isoprene units, i.e. terpenes (carotenes [C12P 23/00](#))}
- 5/02 . acyclic {([C12P 5/007](#) takes precedence)}
- 5/023 . . {Methane}
- 5/026 . . {Unsaturated compounds, i.e. alkenes, alkynes or allenes}

**7/00 Preparation of oxygen-containing organic compounds**

- 7/02 . containing a hydroxy group
- 7/04 . . acyclic
- 7/06 . . . Ethanol, i.e. non-beverage
- 7/065 . . . . {with microorganisms other than yeasts}
- 7/08 . . . . produced as by-product or from waste or cellulosic material substrate
- 7/10 . . . . . substrate containing cellulosic material

- 7/12 . . . . . substrate containing sulfite waste liquor or citrus waste
- 7/14 . . . . . Multiple stages of fermentation; Multiple types of microorganisms or re-use of microorganisms
- 7/16 . . . . . Butanols
- 7/18 . . . . . polyhydric
- 7/20 . . . . . Glycerol
- 7/22 . . . . . aromatic
- 7/24 . . . . . containing a carbonyl group
- 7/26 . . . . . Ketones
- 7/28 . . . . . Acetone-containing products
- 7/30 . . . . . produced from substrate containing inorganic compounds other than water
- 7/32 . . . . . produced from substrate containing inorganic nitrogen source
- 7/34 . . . . . produced from substrate containing protein as nitrogen source
- 7/36 . . . . . produced from substrate containing grain or cereal material
- 7/38 . . . . . Cyclopentanone- or cyclopentadione-containing products
- 7/40 . . . . . containing a carboxyl group {including Peroxycarboxylic acids}
- 7/42 . . . . . Hydroxy-carboxylic acids
- 7/44 . . . . . Polycarboxylic acids
- 7/46 . . . . . Dicarboxylic acids having four or less carbon atoms, e.g. fumaric acid, maleic acid
- 7/48 . . . . . Tricarboxylic acids, e.g. citric acid
- 7/50 . . . . . having keto groups, e.g. 2-ketoglutaric acid
- 7/52 . . . . . Propionic acid; Butyric acids
- 7/54 . . . . . Acetic acid (vinegar C12J)
- 7/56 . . . . . Lactic acid
- 7/58 . . . . . Aldonic, ketoaldonic or saccharic acids (uronic acids C12P 19/00)
- 7/60 . . . . . 2-Ketogulonic acid
- 7/62 . . . . . Carboxylic acid esters

**WARNING**

Groups [C12P 7/62](#) and [C12P 7/625](#) are incomplete pending reclassification of documents from group [C12P 7/6436](#).

Groups [C12P 7/6436](#), [C12P 7/62](#) and [C12P 7/625](#) should be considered in order to perform a complete search.

- 7/625 . . . . . Polyesters of hydroxy carboxylic acids
- 7/64 . . . . . Fats; Fatty oils; Ester-type waxes; Higher fatty acids, i.e. having at least seven carbon atoms in an unbroken chain bound to a carboxyl group; Oxidised oils or fats
- 7/6409 . . . . . Fatty acids
- 7/6418 . . . . . by hydrolysis of fatty acid esters
- 7/6427 . . . . . Polyunsaturated fatty acids [PUFA], i.e. having two or more double bonds in their backbone

**WARNING**

Group [C12P 7/6427](#) is impacted by reclassification into groups [C12P 7/6431](#), [C12P 7/6432](#) and [C12P 7/6434](#).

All groups listed in this Warning should be considered in order to perform a complete search.

- 7/6431 . . . . . Linoleic acids [18:2[n-6]]

**WARNING**

Group [C12P 7/6431](#) is incomplete pending reclassification of documents from group [C12P 7/6427](#).

Groups [C12P 7/6431](#) and [C12P 7/6427](#) should be considered in order to perform a complete search.

- 7/6432 . . . . . Eicosapentaenoic acids [EPA]

**WARNING**

Group [C12P 7/6432](#) is incomplete pending reclassification of documents from group [C12P 7/6427](#).

Groups [C12P 7/6432](#) and [C12P 7/6427](#) should be considered in order to perform a complete search.

- 7/6434 . . . . . Docosahexenoic acids [DHA]

**WARNING**

Group [C12P 7/6434](#) is incomplete pending reclassification of documents from group [C12P 7/6427](#).

Groups [C12P 7/6434](#) and [C12P 7/6427](#) should be considered in order to perform a complete search.

- 7/6436 . . . . . Fatty acid esters

**WARNING**

Group [C12P 7/6436](#) is impacted by reclassification into groups [C12P 7/62](#) and [C12P 7/625](#).

All groups listed in this Warning should be considered in order to perform a complete search.

- 7/6445 . . . . . Glycerides

**WARNING**

Group [C12P 7/6445](#) is impacted by reclassification into group [C12P 7/6458](#).

Groups [C12P 7/6445](#) and [C12P 7/6458](#) should be considered in order to perform a complete search.

- 7/6454 . . . . . by esterification

**WARNING**

Group [C12P 7/6454](#) is impacted by reclassification into group [C12P 7/6458](#).

Groups [C12P 7/6454](#) and [C12P 7/6458](#) should be considered in order to perform a complete search.

7/6458	. . . . by transesterification, e.g. interesterification, ester interchange, alcoholysis or acidolysis	13/02	. Amides, e.g. chloramphenicol {or polyamides; Imides or polyimides; Urethanes, i.e. compounds comprising N-C=O structural element or polyurethanes (peptides C12P 21/00 or C07K)}
	<b>WARNING</b>	13/04	. Alpha- or beta- amino acids {(other amino acids C12P 13/005)}
	Group C12P 7/6458 is incomplete pending reclassification of documents from groups C12P 7/6445, C12P 7/6454, C12P 7/6472, C12P 7/6481 and C12P 7/649.	13/06	. . Alanine; Leucine; Isoleucine; Serine; Homoserine
	All groups listed in this Warning should be considered in order to perform a complete search.	13/08	. . Lysine; Diaminopimelic acid; Threonine; Valine
7/6463	. . . . obtained from glyceride producing microorganisms, e.g. single cell oil	13/10	. . Citrulline; Arginine; Ornithine
7/6472	. . . . containing polyunsaturated fatty acid [PUFA] residues, i.e. having two or more double bonds in their backbone	13/12	. . Methionine; Cysteine; Cystine
	<b>WARNING</b>	13/14	. . Glutamic acid; Glutamine
	Group C12P 7/6472 is impacted by reclassification into group C12P 7/6458.	13/16	. . . using surfactants, fatty acids or fatty acid esters, i.e. having at least seven carbon atoms in an unbroken chain bound to a carboxyl group or a carboxyl ester group
	Groups C12P 7/6472 and C12P 7/6458 should be considered in order to perform a complete search.	13/18	. . . using biotin or its derivatives
7/6481	. . . . Phosphoglycerides (phosphoglycerides having carboxylic acids with less than seven carbon atoms C12P 7/62)	13/20	. . Aspartic acid; Asparagine
	<b>WARNING</b>	13/22	. . Tryptophan; Tyrosine; Phenylalanine; 3,4-Dihydroxyphenylalanine
	Group C12P 7/6481 is impacted by reclassification into group C12P 7/6458.		<b>NOTE</b>
	Groups C12P 7/6481 and C12P 7/6458 should be considered in order to perform a complete search.		Processes for the preparation of different amino acids covered by more than one of the groups C12P 13/222 - C12P 13/227 are classified in group C12P 13/22
7/649	. . . Biodiesel, i.e. fatty acid alkyl esters	13/222	. . . {Phenylalanine}
	<b>WARNING</b>	13/225	. . . {Tyrosine; 3,4-Dihydroxyphenylalanine}
	Group C12P 7/649 is impacted by reclassification into group C12P 7/6458.	13/227	. . . {Tryptophan}
	Groups C12P 7/649 and C12P 7/6458 should be considered in order to perform a complete search.	13/24	. . Proline; Hydroxyproline; Histidine
7/66	. containing the quinoid structure	15/00	<b>Preparation of compounds containing at least three condensed carbocyclic rings {(gibbanes C12P 27/00; naphthacenes C12P 29/00)}</b>
9/00	<b>Preparation of organic compounds containing a metal or atom other than H, N, C, O, S or halogen {(phosphoglycerides, C12P 7/6481)}</b>	17/00	<b>Preparation of heterocyclic carbon compounds with only O, N, S, Se or Te as ring hetero atoms (C12P 13/04 - C12P 13/24 take precedence)</b>
11/00	<b>Preparation of sulfur-containing organic compounds</b>	17/02	. Oxygen as only ring hetero atoms
13/00	<b>Preparation of nitrogen-containing organic compounds</b>	17/04	. . containing a five-membered hetero ring, e.g. griseofulvin {, vitamin C}
13/001	. {Amines; Imines}	17/06	. . containing a six-membered hetero ring, e.g. fluorescein
13/002	. {Nitriles (-CN)}	17/08	. . containing a hetero ring of at least seven ring members, e.g. zearalenone, macrolide aglycons
13/004	. . {Cyanohydrins}	17/10	. Nitrogen as only ring hetero atom
13/005	. {Amino acids other than alpha- or beta amino acids, e.g. gamma amino acids}	17/12	. . containing a six-membered hetero ring
13/007	. {Carnitine; Butyrobetaine; Crotonobetaine}	17/14	. Nitrogen or oxygen as hetero atom and at least one other diverse hetero ring atom in the same ring
13/008	. {containing a N-O bond, e.g. nitro (-NO <sub>2</sub> ), nitroso (-NO)}	17/16	. containing two or more hetero rings {(thiamine open chain analogs C12P 17/167, i.e. not condensed among themselves or through a common carbocyclic ring system)}
		17/162	. . {Heterorings having oxygen atoms as the only ring heteroatoms, e.g. Lasalocid}
		17/165	. . {Heterorings having nitrogen atoms as the only ring heteroatoms}
		17/167	. . {Heterorings having sulfur atoms as ring heteroatoms, e.g. vitamin B1, thiamine nucleus and open chain analogs}
		17/18	. containing at least two hetero rings condensed among themselves or condensed with a common carbocyclic ring system, e.g. rifamycin {(e.g. Rifamycin C12P 17/189)}

- 17/181 . . {Heterocyclic compounds containing oxygen atoms as the only ring heteroatoms in the condensed system, e.g. Salinomycin, Septamycin}
- 17/182 . . {Heterocyclic compounds containing nitrogen atoms as the only ring heteroatoms in the condensed system (alloxazine or isalloxazine, e.g. riboflavine C12P 25/00)}
- 17/183 . . . {containing an indolo[4,3-F,G]quinoline nucleus, e.g. compound containing the lysergic acid nucleus as well as the dimeric ergot nucleus}
- 17/184 . . . {containing a beta-lactam ring, e.g. thienamycin}
- 17/185 . . {Heterocyclic compounds containing sulfur atoms as ring heteroatoms in the condensed system (cebam nucleus C12P 35/00; penam nucleus C12P 37/00)}
- 17/186 . . . {containing a 2-oxo-thieno[3,4-d]imidazol nucleus, e.g. Biotin}
- 17/187 . . . {containing two or more directly linked sulfur atoms, e.g. epithiopiperazines}
- 17/188 . . {Heterocyclic compound containing in the condensed system at least one hetero ring having nitrogen atoms and oxygen atoms as the only ring heteroatoms (ergot-alkaloids C12P 17/183)}
- 17/189 . . . {containing the rifamycin nucleus}
- 19/00 Preparation of compounds containing saccharide radicals (ketoaldonic acids C12P 7/58)**
- NOTE**
- Attention is drawn to the term "saccharide radical" in the first Note following the title of subclass C07H.
- 19/02 . Monosaccharides (2-ketogulonic acid C12P 7/60)
- 19/04 . Polysaccharides, i.e. compounds containing more than five saccharide radicals attached to each other by glycosidic bonds
- 19/06 . . Xanthan, i.e. Xanthomonas-type heteropolysaccharides
- 19/08 . . Dextran
- 19/10 . . Pullulan
- 19/12 . Disaccharides
- 19/14 . produced by the action of a carbohydrase {(EC 3.2.x)}, e.g. by alpha-amylase {, e.g. by cellulase, hemicellulase}
- 19/16 . produced by the action of an alpha-1, 6-glucosidase, e.g. amylose, debranched amylopectin (non-biological hydrolysis of starch C08B 30/00)
- 19/18 . produced by the action of a glycosyl transferase, e.g. alpha-, beta- or gamma-cyclodextrins
- 19/20 . produced by the action of an exo-1,4 alpha-glucosidase, e.g. dextrose
- 19/22 . produced by the action of a beta-amylase, e.g. maltose
- 19/24 . produced by the action of an isomerase, e.g. fructose
- 19/26 . Preparation of nitrogen-containing carbohydrates
- 19/28 . . N-glycosides
- 19/30 . . . Nucleotides
- 19/305 . . . . {Pyrimidine nucleotides}
- 19/32 . . . . having a condensed ring system containing a six-membered ring having two N-atoms in the same ring, e.g. purine nucleotides, nicotineamide-adenine dinucleotide
- 19/34 . . . . Polynucleotides, e.g. nucleic acids, oligoribonucleotides
- 19/36 . . . . Dinucleotides, e.g. nicotineamide-adenine dinucleotide phosphate
- 19/38 . . . Nucleosides
- 19/385 . . . . {Pyrimidine nucleosides}
- 19/40 . . . . having a condensed ring system containing a six-membered ring having two nitrogen atoms in the same ring, e.g. purine nucleosides
- 19/42 . . . Cobalamins, i.e. vitamin B<sub>12</sub>, LLD factor
- 19/44 . Preparation of O-glycosides, e.g. glucosides {(polysaccharides and not substituted disaccharides C12P 19/04, C12P 19/12)}
- 19/445 . . {The saccharide radical is condensed with a heterocyclic radical, e.g. everninomycin, papulacandin}
- 19/46 . . having an oxygen atom of the saccharide radical bound to a cyclohexyl radical, e.g. kasugamycin
- 19/48 . . . the cyclohexyl radical being substituted by two or more nitrogen atoms, e.g. destomycin, neamin
- 19/485 . . . . {Having two saccharide radicals bound through only oxygen to non-adjacent ring carbons of the cyclohexyl radical, e.g. gentamycin, kanamycin, sisomycin, verdamycin, mutamycin, tobramycin, nebramycin, antibiotics 66-40B, 66-40D, XK-62-2, 66-40, G-418, G-52 (see also C12P 19/54)}
- 19/50 . . . . having two saccharide radicals bound through only oxygen to adjacent ring carbon atoms of the cyclohexyl radical, e.g. ambutyrosin, ribostamycin
- 19/52 . . . . . containing three or more saccharide radicals, e.g. neomycin, lividomycin
- 19/54 . . . the cyclohexyl radical being bound directly to a nitrogen atom of two or more  $\begin{array}{c} \text{>N-C-N<} \\ || \\ \text{N} \end{array}$  radicals, e.g. streptomycin
- 19/56 . . having an oxygen atom of the saccharide radical directly bound to a condensed ring system having three or more carbocyclic rings, e.g. daunomycin, adriamycin
- 19/58 . . having an oxygen atom of the saccharide radical directly bound through only acyclic carbon atoms to a non-saccharide heterocyclic ring, e.g. bleomycin, phleomycin
- 19/60 . . having an oxygen of the saccharide radical directly bound to a non-saccharide heterocyclic ring or a condensed ring system containing a non-saccharide heterocyclic ring, e.g. coumermycin, novobiocin {(C12P 19/605)}
- 19/605 . . . {to a 1-benzopyran-2-on (or the chalcones and hydrogenated chalcones thereof, e.g. coumermycin, novobiocin, novenamin)}
- 19/62 . . . the hetero ring having eight or more ring members and only oxygen as ring hetero atoms, e.g. erythromycin, spiramycin, nystatin
- 19/623 . . . . {Avermectin; Milbemycin; Ivermectin; C-076}

19/626	. . . . {Natamycin; Pimaricin; Tenuacetin}	35/00	<b>Preparation of compounds having a 5-thia-1-azabicyclo [4.2.0] octane ring system, e.g. cephalosporin</b>
19/64	. Preparation of S-glycosides, e.g. lincomycin	35/02	. by desacylation of the substituent in the 7 position
<b>21/00</b>	<b>Preparation of peptides or proteins (single cell protein C12N 1/00)</b>	35/04	. by acylation of the substituent in the 7 position
21/005	. {Glycopeptides, glycoproteins}	35/06	. Cephalosporin C; Derivatives thereof
21/02	. having a known sequence of two or more amino acids, e.g. glutathione	35/08	. disubstituted in the 7 position
21/06	. produced by the hydrolysis of a peptide bond, e.g. hydrolysate products (preparing foodstuffs by protein hydrolysis A23J 3/00)	<b>37/00</b>	<b>Preparation of compounds having a 4-thia-1-azabicyclo [3.2.0] heptane ring system, e.g. penicillin</b>
<b>23/00</b>	<b>Preparation of compounds containing a cyclohexene ring having an unsaturated side chain containing at least ten carbon atoms bound by conjugated double bonds, e.g. carotenes (containing heterorings C12P 17/00)</b>	37/02	. in presence of phenylacetic acid or phenylacetamide or their derivatives {not to be used}
<b>25/00</b>	<b>Preparation of compounds containing alloxazine or isalloxazine nucleus, e.g. riboflavin</b>	37/04	. by acylation of the substituent in the 6 position
<b>27/00</b>	<b>Preparation of compounds containing a gibbane ring system, e.g. gibberellin</b>	37/06	. by desacylation of the substituent in the 6 position
<b>29/00</b>	<b>Preparation of compounds containing a naphthacene ring system, e.g. tetracycline (C12P 19/00 takes precedence)</b>	<b>39/00</b>	<b>Processes involving microorganisms of different genera in the same process, simultaneously</b>
<b>31/00</b>	<b>Preparation of compounds containing a five-membered ring having two side-chains in ortho position to each other, and having at least one oxygen atom directly bound to the ring in ortho position to one of the side-chains, one side-chain containing, not directly bound to the ring, a carbon atom having three bonds to hetero atoms with at the most one bond to halogen, and the other side-chain having at least one oxygen atom bound in gamma-position to the ring, e.g. prostaglandins</b>	<b>41/00</b>	<b>Processes using enzymes or microorganisms to separate optical isomers from a racemic mixture</b>
31/005	. {by fermentation or enzyme-using processes from marine organisms, e.g. Plexaura Homomalla}	41/001	. {by metabolizing one of the enantiomers}
<b>33/00</b>	<b>Preparation of steroids</b>	41/002	. {by oxidation/reduction reactions}
	<b>NOTES</b>	41/003	. {by ester formation, lactone formation or the inverse reactions}
	1. Attention is drawn to the definition of steroids in the note following the title of subclass C07J.	41/004	. . {by esterification of alcohol- or thiol groups in the enantiomers or the inverse reaction}
	2. In groups C12P 33/02 - C12P 33/20, the terms "acting", "forming", "hydroxylating", "dehydroxylating" and "dehydrogenating" refer to the action of a microorganism or enzyme rather than other chemical action.	41/005	. . {by esterification of carboxylic acid groups in the enantiomers or the inverse reaction}
33/005	. {Degradation of the lateral chains at position 17}	41/006	. {by reactions involving C-N bonds, e.g. nitriles, amides, hydantoins, carbamates, lactames, transamination reactions, or keto group formation from racemic mixtures}
33/02	. Dehydrogenating; Dehydroxylating	41/007	. . {by reactions involving acyl derivatives of racemic amines}
33/04	. . Forming an aryl ring from A ring	41/008	. . {by reactions involving carbamates}
33/06	. Hydroxylating	41/009	. . {by reactions involving hydantoins or carbamoylamino compounds}
33/08	. . at 11 position	<b>2201/00</b>	<b>Pretreatment of cellulosic or lignocellulosic material for subsequent enzymatic treatment or hydrolysis</b>
33/10	. . . at 11 alpha-position	<b>2203/00</b>	<b>Fermentation products obtained from optionally pretreated or hydrolyzed cellulosic or lignocellulosic material as the carbon source</b>
33/12	. Acting on D ring {(carbons 13 and 14 belong to the C ring; degradation of lateral chains C12P 33/005)}		
33/14	. . Hydroxylating at 16 position		
33/16	. . Acting at 17 position		
33/18	. . . Hydroxylating at 17 position		
33/20	. containing heterocyclic rings {(reactions are also classified in groups C12P 33/00 - C12P 33/18)}		